arXiv:1611.03953v1 [math.AG] 12 Nov 2016

A BIRATIONAL EMBEDDING OF AN ALGEBRAIC CURVE INTO PROJECTIVE PLANE WITH TWO GALOIS POINTS

SATORU FUKASAWA

Abstract. We present a criterion for the existence of a birational embedding of an algebraic curve into projective plane with two Galois points. As an application, we present several new examples of plane curves with two inner Galois points.

1. Introduction

Let C be a (reduced, irreducible) smooth projective curve over an algebraically closed field k of characteristic $p \geq 0$ and let $k(C)$ be its function field. We consider a rational map φ from C to $\mathbb{P}^2$ which is birational onto its image. For a point $P \in \mathbb{P}^2$, if the function field extension $k(\varphi(C))/\pi_P^* k(\mathbb{P}^1)$ induced by the projection π_P is Galois, then P is called a Galois point for $\varphi(C)$. This notion was introduced by H. Yoshihara ([4, 8]). Furthermore, if a Galois point P is a smooth point of $\varphi(C)$ (resp. is contained in $\mathbb{P}^2 \setminus \varphi(C)$), then P is said to be inner (resp. outer). The associated Galois group at P is denoted by G_P. If $\varphi : C \to \varphi(C)$ is isomorphic, then the number of Galois points is completely determined ([2, 8]). However, determining the number in general is difficult. For example, only seven types of plane curves with two inner Galois points are known (see the table in [10]). It is important to find a condition for the existence of two Galois points.

The following proposition appeared in conversations with T. Takahashi [6] and T. Terasoma [7].

Proposition 1. *Let C be a smooth projective curve. Assume that there exist two finite subgroups G_1 and G_2 of the full automorphism group $\mathrm{Aut}(C)$ such that $G_1 \cap G_2 = \{1\}$, and $C/G_i \cong \mathbb{P}^1$ for $i = 1, 2$. Let f, g be generators of function fields of $C/G_1, C/G_2$ respectively. Then, the rational map*

$$\varphi : C \dashrightarrow \mathbb{P}^2;\ (f : g : 1)$$

2010 *Mathematics Subject Classification.* 14H50.

Key words and phrases. Galois point, plane curve, Galois group.

The author was partially supported by JSPS KAKENHI Grant Number 16K05088.

is birational onto its image, and two points $P_1 = (0:1:0)$ *and* $P_2 = (1:0:0)$ *are Galois points for* $\varphi(C)$.

For both points P_1 and P_2 to be inner, or outer, we need additional conditions. In this article, we present a criterion, as follows.

Theorem 1. *Let* C *be a smooth projective curve and let* G_1, G_2 *be different finite subgroups of* $\mathrm{Aut}(C)$. *Then, there exist a morphism* $\varphi : C \to \mathbb{P}^2$ *and different inner Galois points* $\varphi(P_1), \varphi(P_2) \in \varphi(C)$ *such that* φ *is birational onto its image and* $G_{\varphi(P_i)} = G_i$ *for* $i = 1, 2$, *if and only if the following conditions are satisfied.*

(a) $C/G_1 \cong \mathbb{P}^1$ *and* $C/G_2 \cong \mathbb{P}^1$.
(b) $G_1 \cap G_2 = \{1\}$.
(c) *There exist two different points* $P_1, P_2 \in C$ *such that*

$$P_1 + \sum_{\sigma \in G_1} \sigma(P_2) = P_2 + \sum_{\tau \in G_2} \tau(P_1)$$

as divisors.

Remark 1. *For outer Galois points, we have only to replace (c) by*

(c') *There exists a point* $Q \in C$ *such that* $\sum_{\sigma \in G_1} \sigma(Q) = \sum_{\tau \in G_2} \tau(Q)$ *as divisors.*

As an application, we have the following for rational or elliptic curves.

Theorem 2. *Let* $p \neq 2$. *Then, there exist the following morphisms* $\varphi : \mathbb{P}^1 \to \mathbb{P}^2$ *which are birational onto their images.*

(1) $\deg \varphi(C) = 5$ *and there exist two Galois points* $\varphi(P_1), \varphi(P_2) \in \varphi(C)$ *such that* $G_{\varphi(P_i)} \cong \mathbb{Z}/4\mathbb{Z}$ *for* $i = 1, 2$, *in* $p \neq 3$.
(2) $\deg \varphi(C) = 5$ *and there exist two Galois points* $\varphi(P_1), \varphi(P_2) \in \varphi(C)$ *such that* $G_{\varphi(P_i)} \cong (\mathbb{Z}/2\mathbb{Z})^{\oplus 2}$ *for* $i = 1, 2$.
(3) $\deg \varphi(C) = 5$ *and there exist two Galois points* $\varphi(P_1), \varphi(P_2) \in \varphi(C)$ *such that* $G_{\varphi(P_1)} \cong \mathbb{Z}/4\mathbb{Z}$ *and* $G_{\varphi(P_2)} \cong (\mathbb{Z}/2\mathbb{Z})^{\oplus 2}$.
(4) $\deg \varphi(C) = 6$ *and there exist two Galois points* $\varphi(P_1), \varphi(P_2) \in \varphi(C)$ *such that* $G_{\varphi(P_i)} \cong \mathbb{Z}/5\mathbb{Z}$ *for* $i = 1, 2$.

Theorem 3. *Let* $p \neq 3$ *and let* $E \subset \mathbb{P}^2$ *be the curve defined by* $X^3 + Y^3 + Z^3 = 0$. *Then, there exists a morphism* $\varphi : E \to \mathbb{P}^2$ *such that* φ *is birational onto its image,* $\deg \varphi(E) = 4$, *and there exist two inner Galois points for* $\varphi(E)$.

2. Proof of main theorem

Proof of Proposition 1. Let $P_1 = (0 : 1 : 0)$ and $P_2 = (1 : 0 : 0)$. Then, the projection π_{P_1} (resp. π_{P_2}) is given by $(x : y : 1) \mapsto (x : 1)$ (resp. $(x : y : 1) \mapsto (y : 1)$), and hence, $\pi_{P_1} \circ \varphi = (f : 1)$, $\pi_{P_2} \circ \varphi = (g : 1)$. We have only to show that $k(C) = k(f, g)$. Since $k(C)/k(f)$ is Galois, there exist a subgroup H_1 of G_1 such that $H_1 = \mathrm{Gal}(k(C)/k(f, g))$. Similarly, there exists a subgroup H_2 of G_2 such that $H_2 = \mathrm{Gal}(k(C)/k(f, g))$. Since $G_1 \cap G_2 = \{1\}$, we have $H_1 = H_2 = \{1\}$. Therefore, we have $k(C) = k(f, g)$. □

Proof of Theorem 1. We consider the only-if part. Let $\varphi(P_1), \varphi(P_2) \in \varphi(C)$ be inner Galois points such that $G_{\varphi(P_i)} = G_i$ for $i = 1, 2$. Assertion (a) is obvious. The proof of assertion (b) is similar to [1, Lemma 7]. Let D be the divisor induced by the intersection of C and the line $\overline{P_1P_2}$, where $\overline{P_1P_2}$ is the line passing through P_1 and P_2. We can consider the line $\overline{P_1P_2}$ as a point in the images of $\pi_{P_1} \circ \varphi$ and $\pi_{P_2} \circ \varphi$. Since $\pi_{P_1} \circ \varphi$ (resp. $\pi_2 \circ \varphi$) is a Galois covering and $P_2 \in C \cap \overline{P_1P_2}$ (resp. $P_1 \in C \cap \overline{P_1P_2}$), we have

$$(\pi_{P_1} \circ \varphi)^*(\overline{P_1P_2}) = \sum_{\sigma \in G_1} \sigma(P_2), \ (\pi_{P_2} \circ \varphi)^*(\overline{P_1P_2}) = \sum_{\tau \in G_2} \tau(P_1)$$

as divisors (see, for example, [5, III.7.1, III.7.2, III.8.2]). On the other hand, $(\pi_{P_1} \circ \varphi)^*(\overline{P_1P_2}) = D - P_1$ (resp. $(\pi_{P_2} \circ \varphi)^*(\overline{P_1P_2}) = D - P_2$). Therefore, we have

$$D = P_1 + \sum_{\sigma \in G_1} \sigma(P_2) = P_2 + \sum_{\tau \in G_2} \tau(P_1),$$

which is nothing but assertion (c).

We consider the if-part. Let D be the divisor

$$D = P_1 + \sum_{\sigma \in G_1} \sigma(P_2) = P_2 + \sum_{\tau \in G_2} \tau(P_1),$$

by (c). Let $f, g \in k(C)$ be generators of $k(C/G_1)$, $k(C/G_2)$ such that $(f)_\infty = D - P_1$ (i.e., $f \in \mathcal{L}(D - P_1)$), $(g)_\infty = D - P_2$, by (a), where $(f)_\infty$ is the pole divisor of f. Then, $f, g \in \mathcal{L}(D)$. Let $\varphi : C \to \mathbb{P}^2$ be given by $(f : g : 1)$. Similar to Proposition 1, by (b), φ is birational onto its image. The sublinear system of $|D|$ corresponding to $\langle f, g, 1\rangle$ is base-point-free, since $\mathrm{supp}(D) \cap \mathrm{supp}((f) + D) = \{P_1\}$ and $\mathrm{supp}(D) \cap \mathrm{supp}((g) + D) = \{P_2\}$. Therefore, $\deg \varphi(C) = \deg D$, and the morphism $(f : 1)$ (resp. $(g : 1)$) coincides with the projection from the smooth point $\varphi(P_1) \in \varphi(C)$ (resp. $\varphi(P_2) \in \varphi(C)$). □

3. Applications

First, we consider rational curves.

Proof of Theorem 2. We consider (1). Let $\sigma, \tau \in \mathrm{Aut}(\mathbb{P}^1)$ be represented by

$$\begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ -\frac{1}{2} & 1 \end{pmatrix}$$

respectively, by the assumption $p \neq 2$. Let $G_1 = \langle \sigma \rangle$, $G_2 = \langle \tau \rangle$, $P_1 = (2:1)$ and let $P_2 = (-1:1)$. By $p \neq 3$, $P_1 \neq P_2$. Assertion (a) in Theorem 1 is obvious. Note that

$$\{\sigma^i(P_2) | i = 1,2,3\} = \{(1:0),(1:1),(0:1)\} = \{\tau^i(P_1) | i = 1,2,3\}.$$

We have assertion (c) in Theorem 1. Furthermore, we have $\sigma^4 = 1$ and $\tau^4 = 1$. We prove (b) in Theorem 1. Assume by contradiction that $\sigma^i = \tau^j$ for some i, j. If $i = 1$ or 3, then there exists an integer l such that $(\sigma^i)^l = \sigma$. Then, $\tau^{jl}(0:1) = \sigma(0:1) = (-1:1)$. However, there exists no integer i such that $\tau^i(0:1) = (-1:1)$. This is a contradiction. Therefore, we have $i = 2$ and $j = 2$. However, $\sigma^2(1:0) = (0:1) \neq (1:1) = \tau^2(1:0)$.

We consider (2). Let $\alpha \neq 0, 1, -1$, and let $\sigma_\alpha, \tau_\alpha \in \mathrm{Aut}(\mathbb{P}^1)$ be represented by

$$\begin{pmatrix} 0 & 1 \\ \alpha & 0 \end{pmatrix}, \begin{pmatrix} 1 & -\frac{1}{\alpha} \\ 1 & -1 \end{pmatrix},$$

respectively. Let $G_\alpha = \langle \sigma_\alpha, \tau_\alpha \rangle$. Since $\sigma_\alpha \tau_\alpha = \tau_\alpha \sigma_\alpha$, we have $G_\alpha \cong (\mathbb{Z}/2\mathbb{Z})^{\oplus 2}$. We take $\alpha' \neq 0, 1, -1, \alpha$. Let $G_1 = G_\alpha$, $G_2 = G_{\alpha'}$, $P_1 = (1:\alpha')$ and $P_2 = (1:\alpha)$. Note that

$$\{\sigma_\alpha(P_2), \tau_\alpha(P_2), \sigma_\alpha\tau_\alpha(P_2)\} = \{(1:1),(0:1),(1:0)\} = \{\sigma_{\alpha'}(P_1), \tau_{\alpha'}(P_1), \sigma_{\alpha'}\tau_{\alpha'}(P_1)\}.$$

We have assertion (c) in Theorem 1. Assertions (a)(b) are obvious in this case.

We consider (3). We take σ as in (1) and $\sigma_\alpha, \tau_\alpha$ as in (2) Let $P_1 = (1:\alpha)$, $P_2 = (1:-1)$. Note that

$$\{\sigma^i(P_2) | i = 1,2,3\} = \{(1:0),(1:1),(0:1)\} = \{\sigma_\alpha(P_1), \tau_\alpha(P_1), \sigma_\alpha\tau_\alpha(P_1)\}.$$

Furthermore, we have $\sigma^2 \notin \langle \sigma_\alpha, \tau_\alpha \rangle$. Similar to above, we have the assertion by Theorem 1.

We consider (4). Let $\alpha^2 + \alpha - 1 = 0$ and let $\sigma, \tau \in \mathrm{Aut}(\mathbb{P}^1)$ be represented by

$$\begin{pmatrix} 1 & -1 \\ 1 & -\alpha \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ \alpha - 1 & 1 \end{pmatrix}$$

respectively. Let $G_1 = \langle\sigma\rangle$, $G_2 = \langle\tau\rangle$, $P_1 = (\alpha : 2\alpha - 1)$ and let $P_2 = (1 : 1 + \alpha)$. By $p \neq 2$, $P_1 \neq P_2$. Note that

$$\{\sigma^i(P_2) | 1 \leq i \leq 4\} = \{(1:0), (0:1), (1:1), (1:\alpha)\} = \{\tau^i(P_1) | 1 \leq i \leq 4\}.$$

We have assertion (c) in Theorem 1. Furthermore, we have $\sigma^5 = 1$ and $\tau^5 = 1$. Assertions (a)(b) are obvious in this case. □

Remark 2. *For $\sigma \in \mathrm{Aut}(\mathbb{P}^1)$ as in the proof of Theorem 2(1), we can show that the rational function $\sum_{i=0}^{3} \sigma^*(t) \in k(\mathbb{P}^1) = k(t)$ is a generator of $k(\mathbb{P}^1/\langle\sigma\rangle)$. Then, the birational embedding $\varphi : \mathbb{P}^1 \to \mathbb{P}^2$ is represented by*

$$(2(t^4 - 6t^2 + 1)(2t - 1) : (4t^4 - 12t^2 + 8t - 1)(t + 1) : 2t(t + 1)(t - 1)(2t - 1)).$$

Similarly, for Theorem 2(2), we have a birational embedding

$$((t^2 - \alpha)^2(t - \alpha') : (t^2 - \alpha')^2(t - \alpha) : t(t - 1)(t - \alpha)(t - \alpha')).$$

Remark 3. *According to* [3, Theorem 1]*, if $p = 0$, $C = \mathbb{P}^1$ and $\deg \varphi(C) = 6$, then the number of inner Galois points is bounded by two. Our curve in Theorem 2(4) attains this bound.*

Next, we consider elliptic curves. Let $p \neq 3$. Note that if an elliptic curve E admits a Galois covering over $\mathbb{P}^1$ of degree three, then E has an embedding $\varphi : E \to \mathbb{P}^2$ such that the image is the Fermat cubic. To consider the case where $\deg \varphi(E) = 4$, we assume that $E \subset \mathbb{P}^2$ is the curve defined by $X^3 + Y^3 + Z^3 = 0$.

Proof of Theorem 3. Let σ be the automorphism of E given by $(X : Y : Z) \mapsto (\omega X : Y : Z)$, where $\omega^2 + \omega + 1 = 0$. Then, σ is of order three and $E/\langle\sigma\rangle \cong \mathbb{P}^1$. We take a point $Q \in E \setminus \{YZ = 0\}$ such that $\sigma(Q) \neq Q$ and $\sigma^2(Q) \neq Q$. Note that there exists an involution η such that $\eta(Q) = \sigma(Q)$, by the linear system $|Q + \sigma(Q)|$. We take $\tau := \eta\sigma^2\eta$. Then, $\tau(Q) = \sigma(Q)$, τ is of order three and $E/\langle\tau\rangle \cong \mathbb{P}^1$. Let $G_1 = \langle\sigma\rangle$, $G_2 = \langle\tau\rangle$. Then, we have (a) in Theorem 1 for G_1, G_2. Furthermore, we take $P_1 = \tau^2(Q)$ and $P_2 = \sigma^2(Q)$. To prove (b)(c) in Theorem 1, we have only to show that $\sigma^2(Q) \neq \tau^2(Q)$.

Let $R_1 = (1 : 0 : 0)$, $R_2 = (0 : 1 : 0)$ and $R_3 = (0 : 0 : 1)$. Assume by contradiction that $\tau^2(Q) = \sigma^2(Q)$. Then, we have $\eta(\sigma^2(Q)) = \sigma^2(Q)$. For the divisor $D = Q + \sigma(Q) + \sigma^2(Q)$, $\eta^*(D) = D$. Since D is given by $E \cap \overline{R_1Q}$ and the linear system $|D|$ is complete, where $\overline{R_1Q}$ is the line passing through R_1 and Q, η is the restriction of some linear transformation $\hat{\eta} : \mathbb{P}^2 \cong \mathbb{P}^2$. It is well known that

the set of outer Galois points for E is equal to $\Delta = \{R_1, R_2, R_3\}$ (see, for example, [9]). Since $\hat{\eta}(\Delta) = \Delta$ and $\hat{\eta}(\overline{R_1Q}) = \overline{R_1Q} \not\ni R_2, R_3$, we have $\hat{\eta}(R_1) = R_1$ and $\hat{\eta}(\overline{R_2R_3}) = \overline{R_2R_3}$. Then, $\hat{\eta}$ fixes points R_1, Q and the point given by $\overline{R_1Q} \cap \overline{R_2R_3}$. This implies that $\hat{\eta}$ is identity on the line $\overline{R_1Q}$. This is a contradiction.

By Theorem 1, we have the assertion. □

Remark 4. *According to* [3, Theorem 1]*, if* $p = 0$ *and* $\deg \varphi(C) = 4$*, then the number of inner Galois points is bounded by four. It is not known whether the bound is sharp or not, when* C *is an elliptic curve.*

Acknowledgements

The author is grateful to Professor Takeshi Takahashi and Professor Tomohide Terasoma for helpful conversations.

References

[1] S. Fukasawa, Classification of plane curves with infinitely many Galois points, J. Math. Soc. Japan **63** (2011), 195–209.

[2] S. Fukasawa, Complete determination of the number of Galois points for a smooth plane curve, Rend. Sem. Mat. Univ. Padova **129** (2013), 93–113.

[3] K. Miura, Galois points on singular plane quartic curves, J. Algebra **287** (2005), 283–293.

[4] K. Miura and H. Yoshihara, Field theory for function fields of plane quartic curves, J. Algebra **226** (2000), 283–294.

[5] H. Stichtenoth, *Algebraic function fields and codes*, Universitext, Springer-Verlag, Berlin (1993).

[6] T. Takahashi, private communications, July 2012 and September 2015.

[7] T. Terasoma, private communications, September 2015.

[8] H. Yoshihara, Function field theory of plane curves by dual curves, J. Algebra **239** (2001), 340–355.

[9] H. Yoshihara, Galois points for plane rational curves, Far east J. Math. **25** (2007), 273–284; Errata, ibid. **29** (2008), 209–212.

[10] H. Yoshihara and S. Fukasawa, List of problems, available at: http://hyoshihara.web.fc2.com/openquestion.html

Department of Mathematical Sciences, Faculty of Science, Yamagata University, Kojirakawa-machi 1-4-12, Yamagata 990-8560, Japan

E-mail address: s.fukasawa@sci.kj.yamagata-u.ac.jp